

# Algebra Preliminary Examination

August 2021

Attempt all questions, and justify each answer.

## Part I

1. Let  $G$  be a group. Recall that the *commutator subgroup*  $[G, G]$  of  $G$  is the subgroup generated by all commutators  $[g_1, g_2] = g_1^{-1} g_2^{-1} g_1 g_2$  ( $g_1, g_2 \in G$ ). Also recall that a subgroup  $H$  of  $G$  is *characteristic in  $G$* , written  $H \text{ char } G$ , if each automorphism of  $G$  maps  $H$  onto itself.

(a) Define subgroups  $G^{(n)}$  ( $n \in \mathbb{Z}, n \geq 0$ ) inductively as follows:

$$G^{(0)} = G, \quad G^{(n+1)} = [G^{(n)}, G^{(n)}].$$

Prove that  $G^{(n)} \text{ char } G$  for all  $n \geq 0$ .

(b) Suppose that  $G$  is a non-trivial finite group, such that  $G^{(n)} = 1$  for some  $n > 0$ . Prove that  $G$  has a non-trivial characteristic subgroup of prime power order. (*Hint*: consider the subgroup  $G^{(n-1)}$ , where  $n$  is the smallest integer for which  $G^{(n)} = 1$ .)

2. The *holomorph* of a group  $G$ , denoted  $\text{Hol}(G)$ , is defined to be the semidirect product  $G \rtimes_{\phi} \text{Aut}(G)$ , where  $\phi : \text{Aut}(G) \rightarrow \text{Aut}(G)$  is the identity map. Thus we may identify  $\text{Aut}(G)$  with the subgroup  $K = \{(1, \sigma) : \sigma \in \text{Aut}(G)\}$  of the semidirect product  $\text{Hol}(G)$ . As usual we identify  $G$  with the (normal) subgroup  $\{(g, 1) : g \in G\}$  of  $\text{Hol}(G)$ .

Let  $G = \{1, z_1, z_2, z_3\}$  be the non-cyclic group of order 4 (*i.e.*  $G$  is isomorphic to  $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$ ). Prove that  $\text{Hol}(G)$  is isomorphic to the symmetric group  $S_4$ . (*Hint*: Consider the action by left multiplication of  $\text{Hol}(G)$  on the four left cosets  $K, z_1K, z_2K, z_3K$  of  $K$ .)

## Part II

1. Let  $R$  be an integral domain with the property that every ideal generated by two elements of  $R$  is principal.

(a) Prove that every finitely generated ideal of  $R$  is principal.

(b) Suppose that  $R$  also satisfies the ascending chain condition on principal ideals, *i.e.* given any chain of principal ideals  $I_1 \subseteq I_2 \subseteq I_3 \subseteq \dots$ , there exists a positive integer  $k$  such that  $I_k = I_{k+n}$  for all positive integers  $n$ . Prove that  $R$  is a principal ideal domain.

2. Recall that an element  $e$  of a ring  $R$  is *idempotent* if  $e^2 = e$ . In this question all rings are assumed to be commutative and with  $1 \neq 0$ .

(a) Let  $R$  be a ring containing an idempotent  $e$  distinct from  $0, 1$ . Prove that  $R$  is isomorphic to a direct product of two rings. (*Hint*: if  $e$  is idempotent, then so is  $1 - e$ .)

(b) Suppose that  $R$  is a finite ring and that every element of  $R$  is idempotent. Prove that  $R$  is isomorphic to the direct product of finitely many copies of the field with two elements.

**Part III** In this part, all  $R$ -modules  $M$  are assumed to be unital, i.e.  $1.m = m$  for all  $m \in M$ .

1. Recall that given left  $R$ -modules  $D, M, N$ , an  $R$ -module homomorphism  $\phi : M \rightarrow N$  induces a homomorphism of Abelian groups  $\phi' : \text{Hom}_R(D, M) \rightarrow \text{Hom}_R(D, N)$  given by  $\phi'(\alpha) = \phi \circ \alpha$ .

Let  $R$  be a ring with  $1 \neq 0$  and let  $D, L, M, N$  be left  $R$ -modules. Prove that if the sequence

$$0 \rightarrow L \xrightarrow{\phi} M \xrightarrow{\psi} N \rightarrow 0$$

of module homomorphisms is exact, then the sequence of induced homomorphisms of Abelian groups

$$0 \rightarrow \text{Hom}_R(D, L) \xrightarrow{\phi'} \text{Hom}_R(D, M) \xrightarrow{\psi'} \text{Hom}_R(D, N)$$

is also exact.

2. Let  $I = (2, x)$  be the ideal generated by 2 and  $x$  in the ring  $R = \mathbb{Z}[x]$ ,  $x$  being an indeterminate. The ring  $R/I \cong \mathbb{Z}/2\mathbb{Z}$  inherits from  $R$  a natural  $R$ -module structure, with annihilator  $I$ .
- (a) Show that there is an  $R$ -module homomorphism from  $I \otimes_R I$  to  $\mathbb{Z}/2\mathbb{Z}$  mapping  $p(x) \otimes q(x)$  to  $\frac{p(0)}{2} q'(0)$ , where  $q'$  denotes the usual polynomial derivative of  $q$ .
- (b) Show that  $2 \otimes x \neq x \otimes 2$  in  $I \otimes_R I$ .

**Part IV** In this part,  $x$  denotes an indeterminate.

1. This question concerns the polynomial  $f(x) := x^{p^n} - x + 1 \in \mathbb{F}_p[x]$  ( $n \geq 1$ ). We take some fixed algebraic closure  $\mathcal{A}$  of  $\mathbb{F}_p$ , and denote by  $\mathbb{F}_{p^k}$  the unique field of order  $p^k$  contained in  $\mathcal{A}$ . You may assume that each extension of finite degree of  $\mathbb{F}_p$  is Galois over  $\mathbb{F}_p$ , with cyclic Galois group generated by the Frobenius automorphism  $\phi : a \mapsto a^p$ .
- (a) Let  $E$  be the splitting field over  $\mathbb{F}_p$  of  $f(x) = x^{p^n} - x + 1$  in  $\mathcal{A}$ . Show that  $E$  contains  $\mathbb{F}_{p^n}$  as a subfield. (*Hint: If  $\alpha$  is a root of  $f(x)$ , then so is  $\alpha + a$  for each  $a \in \mathbb{F}_{p^n}$ .*)
- (b) Determine the order of the Frobenius automorphism  $\phi : E \rightarrow E$ ,  $\phi : \beta \mapsto \beta^p$ . (*Hint: First compute  $\phi^n(\alpha)$ , where  $\alpha$  is a root of  $f(x)$ .*)
- (c) Show that if  $f(x)$  is irreducible over  $\mathbb{F}_p$ , then  $pn = p^n$ .  
*[Observation (you may omit the easy proof): from  $pn = p^n$  it follows that  $n = 1$  or  $n = p = 2$ .]*
2. Determine the Galois group over  $\mathbb{Q}$  of  $x^4 + 9$ , describing how each automorphism permutes the roots of this polynomial.

# Algebra Preliminary Examination

January 2021

Attempt all questions, and justify each answer.

## Part I

1. Let  $p$  be a prime, and let  $S_p$  denote the symmetric group of degree  $p$ . Prove that if  $P$  is a subgroup of  $S_p$  of order  $p$ , then the normalizer of  $P$  in  $S_p$  has order  $p(p-1)$ .
2. Classify, up to isomorphism, the groups of order 63.

## Part II

1. A *local ring* is a commutative ring with  $1 \neq 0$  that has a unique maximal ideal. Prove that if  $R$  is a local ring with maximal ideal  $M$ , then every element of  $R \setminus M$  is a unit. Also prove that if  $R$  is a commutative ring with  $1 \neq 0$ , in which the set of nonunits forms an ideal  $M$ , then  $R$  is a local ring with maximal ideal  $M$ .
2. Let  $p \in \mathbb{Z}_+$  be prime, and let  $\mathbb{Z}[i]$  denote the usual ring of Gaussian integers  $\{a+bi \mid a, b \in \mathbb{Z}\}$ . For which  $p$  is the quotient ring  $\mathbb{Z}[i]/(p)$  (i) a field, (ii) a product of fields? Justify your answer. (You may use the following facts: (i)  $\mathbb{Z}[i]$  is a Euclidean Domain with respect to the field norm, hence is also a Unique Factorization Domain, and (ii) a prime  $p \in \mathbb{Z}_+$  with  $p \equiv 1 \pmod{4}$  can be written as the sum of two integer squares.)

*Hint:* Use the Chinese Remainder Theorem where appropriate. Also note that a product of fields cannot contain a nonzero nilpotent element.

## Part III

1. Let  $V$  be a finite dimensional vector space over a field  $F$ , and let  $v_1, v_2$  be nonzero elements of  $V$ . Prove that  $v_1 \otimes v_2 = v_2 \otimes v_1$  in  $V \otimes_F V$  if and only if  $v_1 = \lambda v_2$  for some  $\lambda \in F$ .
2. Let  $R$  be a ring with  $1 \neq 0$ , let  $P, M, N$  be  $R$ -modules, and let there be an exact sequence of  $R$ -module homomorphisms  $M \xrightarrow{\phi} N \rightarrow 0$ .
  - (a) Prove that if  $P$  is a direct summand of a free  $R$ -module, then the induced sequence of Abelian group homomorphisms

$$\mathrm{Hom}_R(P, M) \xrightarrow{\phi'} \mathrm{Hom}_R(P, N) \rightarrow 0$$

is exact. (Here  $\phi'$  is the homomorphism  $\psi \mapsto \phi \circ \psi$ .)

- (b) Show by means of an example that in general the induced sequence  $\mathrm{Hom}_R(P, M) \xrightarrow{\phi'} \mathrm{Hom}_R(P, N) \rightarrow 0$  need not be exact.

*Note:* For this question do not assume any result concerning projective modules.

**Part IV** In this part,  $x$  denotes an indeterminate.

1. This question concerns the splitting field over  $\mathbb{Q}$  of the polynomial  $x^4 - 2x^2 - 2 \in \mathbb{Q}[x]$ .
  - (a) Prove that  $x^4 - 2x^2 - 2$  is irreducible over  $\mathbb{Q}$ , and that its roots in  $\mathbb{C}$  are  $\pm\alpha$ ,  $\pm\beta$ , where  $\alpha = \sqrt{1 + \sqrt{3}}$ ,  $\beta = \sqrt{1 - \sqrt{3}}$ .
  - (b) Prove that  $\mathbb{Q}(\alpha) \neq \mathbb{Q}(\beta)$ , and that  $[\mathbb{Q}(\alpha, \beta) : \mathbb{Q}(\alpha)] = 2$ .
  - (c) Prove that the splitting field of  $x^4 - 2x^2 - 2$  has degree 8 over  $\mathbb{Q}$ , and that the Galois group of this polynomial over  $\mathbb{Q}$  is dihedral of order 8.

*Hint for (c):* The Galois group acts faithfully on the set of roots of the polynomial.

2. Let  $\mathbb{F}_p$  denote the field of order  $p$ , let  $f \in \mathbb{F}_p[x]$  be irreducible over  $\mathbb{F}_p$ , and let  $K$  be a splitting field for  $f$  over  $\mathbb{F}_p$ .

Let  $L$  be an intermediate field, i.e.  $\mathbb{F}_p \subseteq L \subseteq K$ . Prove that the irreducible factors of the polynomial  $f$  in  $L[x]$  all have the same degree.

*Hint:* Here is one approach. Let  $g \in L[x]$  be a factor of  $f$  that is irreducible in  $L[x]$ , and let  $\alpha$  be a root of  $g$  in  $K$ . Consider the relationship between  $[L(\alpha) : L]$  and  $[K : L]$ .

# Algebra Preliminary Examination

August 2020

Attempt all questions, and justify each answer.

## Part I

1. Let  $P$  be a Sylow  $p$ -subgroup of a finite group  $G$ . If  $p$  is the smallest prime dividing  $|G|$  and  $P$  is cyclic, prove that  $N_G(P) = C_G(P)$ . (Recall that  $N_G(P)$ ,  $C_G(P)$  denote the normalizer and centralizer of  $P$  in  $G$ , respectively.)

(Hint: Consider the order of the automorphism group of  $P$  and the action of  $N_G(P)$  on  $P$  by conjugation.)

2. (a) Prove that a group of order 105 contains a cyclic normal subgroup of order 35.  
(b) Prove that, up to isomorphism, there is just one non-Abelian group of order 105.

In parts II, III and IV,  $X$  denotes an indeterminate.

## Part II

1. Let  $R$  be a commutative ring with  $1 \neq 0$ . Recall that  $R$  is *Artinian* if it satisfies the descending chain condition on ideals, i.e. if  $I_1 \supseteq I_2 \supseteq \dots$  is a descending chain of ideals of  $R$ , then there exists  $k \in \mathbb{Z}_+$  such that  $I_m = I_k$  for all  $m > k$ .

Let  $S$  be an arbitrary commutative ring with  $1 \neq 0$ , and let  $J$  denote the Jacobson radical of  $S[X]$ . Prove that  $S[X]/J$  is not Artinian.

2. Let  $R$  be the subring of  $\mathbb{Q}[X]$  consisting of all polynomials whose constant term is an integer.

(a) Prove that  $R$  is an integral domain in which every irreducible element is prime.

(b) Prove that  $R$  is not a Unique Factorization Domain.

(Hint: Consider factorizations of the element  $X$ .)

## Part III

1. Let  $k$  be a field, and let  $R = M_2(k)$  be the ring of  $2 \times 2$  matrices over  $k$ . Let  $P$  be the set of  $2 \times 1$  matrices over  $k$ : then  $P$  is an Abelian group under matrix addition, and left matrix multiplication of elements of  $P$  by elements of  $R$  accords  $P$  the structure of a left  $R$ -module.

Prove that the  $R$ -module  $P$  is projective, but not free.

2. Let  $R = \mathbb{Z}[X]$ , let  $I \subset R$  be the ideal generated by  $2, X$ , and let  $M = I \otimes_R I$ .

Prove that the element  $2 \otimes 2 + X \otimes X \in M$  cannot be written as a simple tensor  $a \otimes b$  ( $a, b \in I$ ).

(Hint: Use a suitable  $R$ -module homomorphism defined on  $M$ .)

#### Part IV

1. Prove that  $\mathbb{Q}(\sqrt{5 + 2\sqrt{5}})$  is a Galois extension of  $\mathbb{Q}$ , and determine its Galois group.
2. Let  $F$  be a field (possibly infinite) of finite characteristic  $p$ , and let  $a \in F$  be an element not of form  $b^p - b$  for any  $b \in F$ . Let  $f = X^p - X - a \in F[X]$ .

(a) Prove that the polynomial  $f$  is separable and irreducible over  $F$ .

(b) Prove that if  $\alpha$  is a root of  $f$  in an extension field of  $F$ , then  $F(\alpha)$  is a splitting field for  $f$  over  $F$ .

(Hint: Consider the effect of substituting  $X + 1$  for  $X$  in the polynomial  $f$ .)

# ALGEBRA PRELIMINARY EXAM

JANUARY 2020

**Instructions:** Attempt *all* problems in all four parts. Justify each answer.

**General Assumptions:** Unless explicitly stated otherwise, all rings have  $1 \neq 0$  [and their subrings contain 1] and all modules are unitary.

## Part I

1. Let  $G$  be a finite group and  $\phi : G \rightarrow H$  a *surjective* homomorphism. Prove that if  $y \in H$  is such that  $|y| = p^r$ , for some prime  $p$  and  $r \in \mathbb{Z}_{>0}$ , then there is  $x \in G$  such that  $\phi(x) = y$  and  $|x| = p^s$ , for some  $s \in \mathbb{Z}_{>0}$ .  
[Hint: Let  $g \in G$  such that  $\phi(g) = y$ , and write  $|g| = n \cdot p^k$ , where  $p \nmid n$ .]
2. Let  $G$  be a group of order 60 and assume that 4 divides  $|Z(G)|$  [where  $Z(G)$  denotes the *center* of  $G$ ]. Prove that  $G$  must be Abelian.

## Part II

1. Let  $I$  be the ideal of  $\mathbb{Z}[x]$  generated by 7 and  $x^2 + 1$ . Prove that  $I$  is a maximal ideal.
2. Let  $R$  be an *integral domain* such that for any descending chain of ideals

$$I_1 \supseteq I_2 \supseteq I_3 \supseteq \cdots$$

there is a positive integer  $N$  such that  $I_i = I_N$  for all  $i \geq N$ . Prove that  $R$  is a field.

## Part III

1. Let  $R$  be a subring of  $S$ . Prove that  $S \otimes_R S \neq 0$ .
2. Let  $R$  be a ring containing  $\mathbb{Z}$  such that  $R$  is a free  $\mathbb{Z}$ -module of finite rank  $n > 0$  and every non-zero ideal of  $R$  has a non-zero element of  $\mathbb{Z}$ . Prove that for every non-zero ideal  $I$  we have that  $R/I$  is finite.

## Part IV

1. Given a prime  $p$  and a positive integer  $n$ , show that there is an *Abelian* extension [i.e., Galois with Abelian Galois group]  $K$  of  $\mathbb{Q}$  with  $[K : \mathbb{Q}] = p^n$ .
2. Let  $F$  be a field of characteristic  $p$  with exactly  $p^r$  elements. If  $K$  is a finite extension of  $F$  with  $K = F[\alpha]$ , for some  $\alpha \in K$ , and  $f$  is the minimal polynomial of  $\alpha$  over  $F$ , then show that if  $\beta$  is another root of  $f$ , then  $\beta \in K$  and  $\beta = \alpha^{p^k}$  for some  $k \in \mathbb{Z}$ .

# ALGEBRA PRELIMINARY EXAM

AUGUST 2019

**Instructions:** Attempt *all* problems in all four parts. Justify each answer.

**General Assumptions:** Unless explicitly stated otherwise, all rings have  $1 \neq 0$  [and their subrings contain 1] and all modules are unitary.

## Part I

1. Let  $G_1, G_2$  be groups,  $N \trianglelefteq G_1$ , and  $\phi : G_1 \rightarrow G_2$  be an onto homomorphism such that  $N \cap \ker(\phi) = \{1\}$ . Prove that for  $x \in N$  we have that  $C_{G_2}(\phi(x)) = \phi(C_{G_1}(x))$ . [Remember:  $C_G(x) \stackrel{\text{def}}{=} \{g \in G : gx = xg\}$  is the *centralizer* of  $x$  in  $G$ .]
2. Let  $G$  be a group of order  $992 = 2^5 \cdot 31$ . Prove that either  $G$  has a normal subgroup of order  $32 = 2^5$  or it has a normal subgroup of order 62.

## Part II

1. Let  $R$  be a UFD with exactly two non-associate prime elements  $p$  and  $q$  [i.e.,  $p$  and  $q$  are non-associate primes and every prime is an associate of either  $p$  or  $q$ ]. Prove that  $R$  is a PID.
2. Let  $R$  be a PID and  $P$  a prime ideal of  $R[x]$  such that  $P \cap R \neq \{0\}$ . Prove that there is  $p \in R$  prime [in  $R$ ] such that either  $P = (p)$  or  $P = (p, f)$  for some  $f$  prime in  $R[x]$ .

## Part III

1. Let  $R$  be a commutative ring and  $M$  an  $R$ -module. Prove that  $R \otimes_R \text{Hom}_R(R \oplus R, M)$  is projective if and only if  $M$  is projective.
2. Let  $R$  be a commutative ring,  $M$  and  $N$  be  $R$ -modules and  $M'$  and  $N'$  be submodules of  $M$  and  $N$  respectively. Define  $L$  as the submodule of  $M \otimes_R N$  generated by the set
$$\{x \otimes y \in M \otimes_R N : x \in M' \text{ or } y \in N'\}.$$
Show that  $M/M' \otimes_R N/N' \cong (M \otimes_R N)/L$ .

## Part IV

1. Let  $F = \mathbb{Q}(\sqrt[3]{2} \cdot \zeta)$ , where  $\zeta = -1/2 + \sqrt{3}i/2$  [a primitive third root of unity]. Prove that  $-1$  is not a sum of squares in  $F$ , i.e., there is no positive integer  $n$  and  $\alpha_1, \dots, \alpha_n \in F$  such that  $-1 = \alpha_1^2 + \dots + \alpha_n^2$ .
2. Let  $F$  be a field of characteristic 0 and  $K/F$  be a field extension of degree  $n$  such that there is a root of unity  $\zeta$  in the algebraic closure of  $K$  such that  $K \subseteq F[\zeta]$ . Prove that if  $d \mid n$ , there is  $\alpha \in K$  such that the minimal polynomial of  $\alpha$  over  $F$  has degree  $d$ .



# ALGEBRA PRELIMINARY EXAM

AUGUST 2018

**Instructions:** Attempt *all* problems in all four parts. Justify your answers.

**General assumptions:** All rings have  $1 \neq 0$ , their subrings contain 1, and all modules are unitary.

## Part I

1. Let  $G$  be a (possibly infinite) group, and suppose that  $G$  contains a subgroup  $H \neq G$  whose index  $[G : H]$  is finite. Prove that  $G$  contains a normal subgroup  $N \neq G$  of finite index.
2. Prove that every group of order 70 contains a cyclic, normal subgroup of order 35.

## Part II

1. Let  $R$  be a commutative ring in which every element is either a unit or nilpotent. Prove that  $R$  has exactly one prime ideal.
2. If  $R$  is an integral domain, prove that there are infinitely many ideals in  $R[x]$  that are both prime and principal.

## Part III

1. Let  $R$  be a ring, possibly non-commutative, and suppose that
$$0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$$
is a short exact sequence of left  $R$ -modules, with  $M'$  and  $M''$  finitely generated. Prove that  $M$  is finitely generated.
2. Let  $M$  be a finitely-generated  $\mathbb{Z}$ -module, and let  $T \subset M$  be its torsion submodule. Show that the torsion submodule of  $M \otimes_{\mathbb{Z}} M$  has at least  $|T|$  elements.

## Part IV

1. Let  $p$  be a prime and suppose that  $f \in \mathbb{F}_p[x]$  is an irreducible polynomial. Let  $K$  be a degree 2 extension of  $\mathbb{F}_p$  and suppose that there exist non-constant polynomials  $g, h \in K[x]$  such that  $f = gh$ . If  $g$  is an irreducible polynomial of degree 5, what is the degree of  $f$ ?
2. Suppose that  $f \in \mathbb{Q}[x]$  is an irreducible degree 4 polynomial, and  $K/\mathbb{Q}$  is an extension such that  $f$  has exactly one root in  $K$ . Let  $G$  be the Galois group of  $f$ , and show that  $|G|$  is divisible by 12.

# ALGEBRA PRELIMINARY EXAM

AUGUST 2017

**Instructions:** Attempt *all* problems in all four parts. Justify your answer.

**General Assumptions:** Unless explicitly stated otherwise, all rings have  $1 \neq 0$  [and their sub-rings contain 1] and all modules are unitary.

## Part I

1. Suppose that  $H$  is a subgroup of a finite group  $G$  of index  $p$ , where  $p$  is the smallest prime dividing the order of  $G$ . Prove that  $H$  is normal in  $G$ .
2. Show that every group of order 222 is solvable.

*Fun fact: The University of Tennessee was established 222 years ago.*

## Part II

1. Let  $I$  and  $J$  be ideals of a ring  $R$  and assume that  $P$  is a prime ideal of  $R$  that contains  $I \cap J$ . Prove that either  $I$  or  $J$  is contained in  $P$ .
2. Let  $R$  be an integral domain and suppose that every prime ideal in  $R$  is principal. Prove that  $R$  is a PID.

## Part III

1. Let  $V$  be a Noetherian right  $R$ -module, and  $\theta : V \rightarrow V$  a homomorphism.
  - (a) Show that  $\ker(\theta^{n+1}) = \ker(\theta^n)$  for some  $n \geq 1$ .
  - (b) If  $\theta$  is onto, show that it is one-to-one.
2. An  $R$ -**projection** is defined to be an  $R$ -module homomorphism  $\varphi : R^n \rightarrow R^n$  such that  $\varphi^2 = \varphi$ . Prove that a finitely generated  $R$ -module  $M$  is projective if and only if it is isomorphic to the image of some  $R$ -projection.

## Part IV

1. Let  $F \subseteq E$  be fields and suppose  $0 \neq \alpha \in E$  with  $E = F(\alpha)$ . Assume that some power of  $\alpha$  lies in  $F$  and let  $n$  be the smallest positive integer such that  $\alpha^n \in F$ .
  - (a) If  $\alpha^m \in F$  with  $m > 0$ , show that  $m$  is a multiple of  $n$ .
  - (b) If  $E$  is a separable extension of  $F$ , prove that the characteristic of  $F$  does not divide  $n$ .
  - (c) If every root of unity of  $E$  lies in  $F$ , show that  $[E : F] = n$ .
2. Let  $F$  be a field of characteristic 0 and let  $E$  be a finite Galois extension of  $F$ .
  - (a) If  $0 \neq \alpha \in E$  with  $E = F(\alpha)$ , show that  $F(\alpha^2) \neq E$  if and only if there exists  $\sigma \in \text{Gal}(E/F)$  with  $\sigma(\alpha) = -\alpha$ .
  - (b) Prove that there exists an element  $\alpha \in E$  with  $E = F(\alpha^2)$ .

ALGEBRA PRELIMINARY EXAMINATION  
SPRING 2017

- Attempt all four parts. Justify your answers.

**Part I.**

1. Show that a group of order 255 is not a simple group.
2. A group  $G$  has a cyclic normal subgroup of order 2016. If  $G$  also has a subgroup of order 2017, then show that  $G$  has a cyclic subgroup of order  $(2016) \times (2017)$ .

**Part II.**

Note: Rings are assumed to be commutative and with  $1 \neq 0$ .

1. Let  $A$  and  $B$  be rings. Show that each ideal of  $A \times B$  is of the form  $I \times J$ , where  $I$  is an ideal of  $A$  and  $J$  is an ideal of  $B$ .
2. Let  $R$  be a ring, let  $X$  be an indeterminate and let  $S := \{X^n \mid 0 \leq n \in \mathbb{Z}\}$ . If  $S^{-1}R[[X]]$  is a field, then show that  $R$  is a field.

**Part III.**

Note: Rings are assumed to be commutative with  $1 \neq 0$  and modules are assumed to be unitary.

1. Let  $A$  be a ring and let  $M, N$  be finitely generated projective (left)  $A$ -modules. Show that  $\text{Hom}_A(M, N)$  is a finitely generated projective  $A$ -module.
2. Let  $R$  be a PID and let  $I, J$  be ideals of  $R$ . If  $I \neq R \neq J$ , then show that  $(R/I) \oplus (R/J)$  and  $(R/I) \otimes_R (R/J)$  are not isomorphic as (left)  $R$ -modules.

**Part IV.**

Note: In what follows,  $X$  is an indeterminate.

1. Let  $K$  be an extension-field of  $\mathbb{Q}$  such that  $K/\mathbb{Q}$  is Galois with Galois group  $\mathbb{Z}_{30}$ . Suppose each of  $f, g \in \mathbb{Q}[X]$  is an irreducible polynomial of degree 6 and  $f$  has a root  $a \in K$ . If  $g$  has a root in  $K$ , then show that  $g$  has all its roots in  $\mathbb{Q}[a]$ .
2. Let  $F \subset K$  be finite fields of characteristic 5 and suppose  $g \in F[x]$  is irreducible in  $F[x]$ . If  $g$  has degree 11, then show that either  $g$  is irreducible in  $K[x]$  or all its roots are in  $K$ .

ALGEBRA PRELIMINARY EXAMINATION  
Fall 2016

- Attempt all four parts. Justify your answers.

**Part I.**

1. Let  $p$  be a prime number and  $G$  be a non-Abelian group of order  $p^3$ . Show that  $G$  has at least 3 (distinct) subgroups of index  $p$ .
2. Let  $G$  be a group of order  $p^3q$ , where  $p, q$  are distinct prime numbers. If no Sylow  $p$ -subgroup of  $G$  is normal and also no Sylow  $q$ -subgroup of  $G$  is normal, then show that  $G$  has order 24.

**Part II.**

Note: Rings are tacitly assumed to be commutative and with  $1 \neq 0$ .

1. Let  $R$  be a ring,  $X$  an indeterminate and  $h : R[X] \rightarrow R[[X]]$  a ring-homomorphism such that  $h(a) = a$  for all  $a \in R$ . Show that  $h$  is not surjective.
2. Let  $R$  be an integral domain with at least 3 (distinct) maximal ideals. Given maximal ideals  $M$  and  $N$  of  $R$ , show that  $R_M \cap R_N \neq R$ . (Here localization of  $R$  at a prime ideal is naturally identified as a ring in between  $R$  and the quotient-field of  $R$ .)

**Part III.**

Note: Rings are assumed to be commutative and with  $1 \neq 0$  and modules are assumed to be unitary.

1. Let  $R$  be a ring and let  $a \in R$  be a nonzero element of  $R$  such that  $a^3 = a$ . Show that the ideal  $Ra$  is a projective  $R$ -module.
2. Let  $R$  be a PID and let  $M$  be a finitely generated  $R$ -module. For a maximal ideal  $Q$  of  $R$ , let  $\delta(Q, M)$  denote the dimension of  $M \otimes_R R/Q$  as a vector-space over the field  $R/Q$ . Let  $\delta(M)$  denote the  $\sup\{\delta(Q, M)\}$ , where the supremum is taken over all maximal ideals  $Q$  of  $R$ . Show that as an  $R$ -module,  $M$  has a generating set of cardinality  $\delta(M)$  and any generating set of  $M$  has cardinality at least  $\delta(M)$ .

**Part IV.**

Note: In what follows,  $X$  is an indeterminate.

1. Let  $f(X)$  be a monic polynomial with rational coefficients. Assume  $f(X)$  is irreducible in  $\mathbb{Q}[X]$  and the Galois-group of  $f(X)$  over  $\mathbb{Q}$  is a group of order 99. What is the degree of  $f(X)$ ?
2. Compute the Galois group of  $X^6 - 9$  over  $\mathbb{Q}$ .

# ALGEBRA PRELIMINARY EXAM

JANUARY 2016

**Instructions:** Attempt *all* problems in all four parts. Justify each answer.

**General Assumptions:** Unless explicitly stated otherwise, all rings have  $1 \neq 0$  [and their subrings contain 1] and all modules are unitary.

## Part I

1. Let  $G$  be a finite group and  $H$  be a subgroup of  $G$ . Prove that  $n_p(H) \leq n_p(G)$ , where  $n_p(X)$  denotes the number of Sylow  $p$ -subgroups of  $X$ .
2. Let  $G$  be a group of order  $p^n$  for some prime  $p$  and positive integer  $n$ . Prove that if  $1 \neq H \trianglelefteq G$ , then  $Z(G) \cap H \neq 1$ . [Here  $Z(G)$  denotes the center of  $G$ .]

## Part II

1. Let  $R$  be a *Boolean ring*, i.e., a ring [with 1] for which  $a^2 = a$  for all  $a \in R$ . [You can use without proof the well known fact that if  $R$  is Boolean, then it is commutative of characteristic 2.]
  - (a) Prove that if  $R$  is finite, then its order is a power of 2.
  - (b) Prove that every prime ideal of  $R$  is maximal.
2. Show that  $R \stackrel{\text{def}}{=} \mathbb{Z}[x_1, x_2, x_3, \dots]/(x_1x_2, x_3x_4, x_5x_6, \dots)$  has infinitely many distinct *minimal* prime ideals. [ $P$  is a minimal prime ideal if it is prime and whenever  $Q \subseteq P$ , with  $Q$  also prime, we have  $Q = P$ .]

## Part III

1. Let  $F$  be a field and  $M$  be a *torsion*  $F[x]$ -module. Prove that if there is  $m_0 \in M$ , with  $m_0 \neq 0$ , and an *irreducible*  $f \in F[x]$  such that  $f \cdot m_0 = 0$ , then  $\text{Ann}(M) \subseteq (f)$ .
2. Let  $R$  be an integral domain and  $I$  a principal ideal of  $R$ . Prove that  $I \otimes_R I$  has no non-zero torsion element [i.e., if  $m \in I \otimes_R I$ , with  $m \neq 0$ , and  $r \in R$  with  $rm = 0$ , then  $r = 0$ ].

## Part IV

1. Let  $K/F$  be an algebraic field extension and  $\text{Emb}(K/F)$  denote the set of field homomorphisms  $\sigma : K \rightarrow \bar{K}$  such that  $\sigma(a) = a$  for all  $a \in F$ . [Here  $\bar{K}$  is a fixed algebraic closure of  $K$ .]
  - (a) Prove that if  $\alpha$  is a root of a [not necessarily irreducible] non-zero polynomial  $f \in F[x]$  with  $\deg(f) = n$ , then  $\text{Emb}(F[\alpha]/F)$  has at most  $n$  elements.
  - (b) Give an example of an algebraic extension  $K/F$  of degree greater than one for which  $\text{Emb}(K/F)$  has a single element.
2. Let  $F = \mathbb{Q}[\sqrt{2}]$  and  $K = \mathbb{Q}[\sqrt[8]{2}, i]$ .
  - (a) Prove that  $K/F$  is Galois with  $[K : F] = 8$ .
  - (b) Prove that  $\text{Gal}(K/F)$  has a non-normal subgroup. [This implies that it is the dihedral group of order 8, as it is the only group of order 8 with this property.]

# ALGEBRA PRELIMINARY EXAM

AUGUST 2015

**Instructions:** Attempt *all* problems in all four parts. Justify each answer.

**General Assumptions:** Unless explicitly stated otherwise, all rings have  $1 \neq 0$  [and their subrings contain 1] and all modules are unitary.

## Part I

1. Let  $G$  be a *non-Abelian* group of order  $p^3$ ,  $[G, G] = \langle xyx^{-1}y^{-1} : x, y \in G \rangle$  be its commutator subgroup and  $Z(G)$  be its center. Show that  $|Z(G)| = p$  and that  $Z(G) = [G, G]$ .
2. Let  $G_1$  and  $G_2$  be groups of order 81 acting *faithfully* [i.e., only 1 acts as the identity function] on sets  $X_1$  and  $X_2$ , respectively, with 9 elements each. Show that there is an isomorphism  $\psi : G_1 \rightarrow G_2$ .

## Part II

1. Let  $D$  be a *finite* division ring. Prove that  $D$  has a prime power number of elements. [Hint: Consider the center  $Z(D) = \{a \in D : ax = xa \text{ for all } x \in D\}$ .]
2. Let  $p \in \mathbb{Z}$  prime and

$$f = a_{2n+1}x^{2n+1} + \cdots + a_1x + a_0 \in \mathbb{Z}[x].$$

Prove that if  $p^3 \nmid a_0$ ,  $p^2 \mid a_0, a_1, \dots, a_n$ ,  $p \mid a_{n+1}, a_{n+2}, \dots, a_{2n}$  and  $p \nmid a_{2n+1}$ , then  $f$  is irreducible in  $\mathbb{Q}[x]$ .

## Part III

1. Let  $R$  be a commutative ring. An  $R$ -module is *Artinian* if it satisfies the *descending chain condition for submodules*. [I.e., if  $S_1 \supseteq S_2 \supseteq S_3 \supseteq \cdots$  is a chain of submodules, then there is a  $i_0$  such that for all  $i \geq i_0$ , we have  $S_i = S_{i_0}$ .] Show that if  $L$  and  $N$  are Artinian  $R$ -modules and we have a short exact sequence

$$0 \longrightarrow L \xrightarrow{\psi} M \xrightarrow{\phi} N \longrightarrow 0,$$

then  $M$  is also Artinian.

2. Let  $R$  be a commutative ring such that every  $R$ -module is free. Prove that  $R$  is a field.

## Part IV

1. Let  $\mathbb{F}_p$  be the field with  $p$  elements, and  $t$  be an indeterminate. Let  $f(t), g(t) \in \mathbb{F}_p[t] \setminus \{0\}$ , with  $\max\{\deg f, \deg g\} < p$  and  $f(t)/g(t) \notin \mathbb{F}_p$ . Show that the extension  $\mathbb{F}_p(t)/\mathbb{F}_p(f(t)/g(t))$  is separable.
2. Suppose that  $f = \prod_{i=1}^N (x - \alpha_i) \in \mathbb{Q}[x]$  [with  $\alpha_i \in \mathbb{C}$ ] is *irreducible* in  $\mathbb{Q}[x]$  and let  $f_n \stackrel{\text{def}}{=} \prod_{i=1}^N (x - \alpha_i^n)$ . Prove that for each  $n$ , there is  $g_n \in \mathbb{Q}[x]$  *irreducible* and a positive integer  $k_n$  such that  $f_n = g_n^{k_n}$ .

ALGEBRA PRELIMINARY EXAMINATION  
Fall 2014

Attempt all four parts. Justify your answers.

Part I.

1. Show that  $S_4$  (the group of permutations of  $\{1, 2, 3, 4\}$ ) does not have a subgroup isomorphic to  $Q_8$  (the quaternion-group of order 8).
2. Let  $G$  be a group of order 2014. Show that  $G$  is cyclic if and only if  $G$  has a normal subgroup of order 2.

Part II.

*Note: Rings are assumed to be commutative and with  $1 \neq 0$ , subrings are assumed to contain 1 and ring-homomorphisms are assumed to map 1 to 1.*

1. Let  $R$  be an integral domain with only finitely many units. Show that the intersection of all maximal ideals of  $R$  is 0.
2. Let  $R$  be a ring such that each non-unit of  $R$  is nilpotent. Let  $X$  be an indeterminate and let  $f \in R[[X]]$ . Show that  $f^n = f$  for some integer  $n \geq 2$  if and only if either  $f = 0$  or  $f^{n-1} = 1$ .

Part III.

*Note: Rings are assumed to be commutative and with  $1 \neq 0$  and modules are assumed to be unitary.*

1. Let  $L$  be a module over a ring  $R$  and let  $M, N$  be  $R$ -submodules of  $L$ . Show that if  $(M + N)/(M \cap N)$  is a projective  $R$ -module then  $M/(M \cap N)$  is also a projective  $R$ -module.
2. Let  $R$  be a PID with infinitely many prime ideals and let  $M$  be a finitely generated  $R$ -module. Show that  $M$  is a torsion  $R$ -module if and only if  $M \otimes_R R/P = 0$  for all but finitely many prime ideals  $P$  of  $R$ .

Part IV.

*Note: In what follows,  $X$  is an indeterminate.*

1. Let  $f(X) := X^5 + 3X^3 + X^2 + 3 \in \mathbb{Q}[X]$ . Let  $K$  be the splitting field of  $f(X)$  over  $\mathbb{Q}$ . Compute  $[K : \mathbb{Q}]$ .
2. Let  $f(X) := X^3 + X + 1 \in \mathbb{Q}[X]$ . Let  $F$  be a finite Galois extension of  $\mathbb{Q}$  such that the Galois group of  $F$  over  $\mathbb{Q}$  is an Abelian group. Show that  $f$  is irreducible in  $F[X]$ .

**Algebra Preliminary Exam      January 2014**

Attempt all problems and justify all your answers. All rings have a  $1 \neq 0$ , all ring homomorphisms send 1 to 1, and all  $R$ -modules are unitary.

**Part I. Groups**

1. Show that every group of order 1,225 is abelian.
2. Let  $n \geq 2$ . Show that there is a nontrivial homomorphism  $f : S_n \rightarrow \mathbb{Z}/n\mathbb{Z}$  (i.e.,  $\ker f \neq S_n$ ) if and only if  $n$  is even.

**Part II. Rings**

1. Let  $R$  be a commutative ring. Show that  $J(R[X]) = \text{nil}(R[X])$ .  
( $J(A)$  and  $\text{nil}(A)$  are the Jacobson and nil radicals of  $A$ .)
2. Let  $R$  be a PID.
  - (a) Show that  $R$  satisfies ACC on ideals.
  - (b) Show that every nonzero prime ideal of  $R$  is maximal.

**Part III. Modules**

1. Let  $R$  be a ring and  $M$  a nonzero  $R$ -module. Show that  $M = A \oplus B$  for proper submodules  $A$  and  $B$  of  $M$  if and only if there is a nonzero, nonidentity homomorphism  $f : M \rightarrow M$  with  $f^2 = f$ .
2. Let  $R$  be a commutative ring,  $I$  a proper ideal of  $R$ , and  $M$  an  $R$ -module. Show that  $(R/I) \otimes_R M$  and  $M/IM$  are isomorphic as  $R$ -modules.

**Part IV. Fields**

1. Let  $K$  a subfield of a field  $F$ . Show that there is a subring of  $F$  containing  $K$  that is a PID, but not a field, if and only if the extension  $F/K$  is not algebraic.
2. Determine the Galois group of  $f(X) = X^{10} + X^8 + X^6 + X^2$  over  $\mathbb{Z}/2\mathbb{Z}$ .



## Algebra Preliminary Exam

August 2013

Attempt all problems and justify all your answers.  
All rings have an identity  $1 \neq 0$ , all ring homomorphisms send 1 to 1, and all  $R$ -modules are unitary.

### Part I.

- (a) Let  $p$  and  $q$  be (not necessarily distinct) prime numbers. Show that a group  $G$  with  $|G| = pq$  is either abelian or  $Z(G) = \{e\}$ .

(b) Give an example of a nonabelian group  $G$  whose order is the product of three (not necessarily distinct) primes and  $Z(G) \neq \{e\}$ .
- (a) Let  $G$  be a group with  $|G| = 100$ . Show that  $G$  is abelian if and only if its Sylow 2-subgroup is normal.

(b) Give an example of a nonabelian group of order 100.

### Part II.

- Let  $R$  and  $S$  be commutative rings with  $1 \neq 0$ . Show that every ideal of  $R \times S$  has the form  $I \times J$  for  $I$  an ideal of  $R$  and  $J$  an ideal of  $S$ .
- Let  $R$  be a commutative ring with  $1 \neq 0$ . Show that  $f(X) = a_0 + a_1X + \cdots + a_nX^n$  is a unit in  $R[X]$  if and only if  $a_0$  is a unit in  $R$  and  $a_1, \dots, a_n$  are nilpotent.

### Part III

1. Let  $P$  and  $Q$  be finitely generated projective  $R$ -modules over a commutative ring  $R$  with  $1 \neq 0$ . Show that  $\text{Hom}_R(P, Q)$  is a finitely generated projective  $R$ -module.
2. Let  $R$  be a commutative ring with  $1 \neq 0$ ,  $S$  a nonempty multiplicatively closed subset of  $R$ , and  $M$  an  $R$ -module. Show that  $(S^{-1}R) \otimes_R M$  and  $S^{-1}M$  are isomorphic as  $S^{-1}R$ -modules.

### Part IV.

1. Let  $p$  and  $q$  be distinct prime numbers,  $F$  a subfield of a field  $K$ , and  $f(X), g(X) \in F[X]$  be irreducible with  $\deg(f(X)) = p$  and  $\deg(g(X)) = q$ . Let  $a, b \in K$  be roots of  $f(x)$  and  $g(X)$ , respectively. Show that  $[F(a, b) : F] = pq$ .
2. (a) Let  $F$  be a splitting field for  $f(X) \in \mathbb{Q}[X]$  over  $\mathbb{Q}$  with abelian Galois group  $G$ . Show that every subfield  $L$  of  $F$  is a splitting field over  $\mathbb{Q}$  for some polynomial  $g(X) \in \mathbb{Q}[X]$ .  
(b) Give an example to show that if  $G$  is not abelian in part (a), then some  $L$  need not be a splitting field.

# ALGEBRA PRELIMINARY EXAM

JANUARY 2013

**Instructions:** Attempt *all* problems in all four parts. Justify each answer.

**General Assumptions:** Unless explicitly stated otherwise, all rings have  $1 \neq 0$  [and their subrings contain 1] and all modules are unitary.

## Part I

1. Let  $p$  and  $q$  be prime numbers such that  $q < p$  and  $q$  does not divide  $p^2 - 1$ . Prove that every group of order  $p^2q$  is Abelian.
2. Let  $G$  be a finite *simple* group. Show that if  $p$  is the *largest* prime dividing  $|G|$ , then there is no subgroup  $H \leq G$  such that  $1 < |G : H| < p$ .

## Part II

1. Let  $R$  be a ring not necessarily having 1 [or commutative], with at least two elements and such that for all non-zero  $a \in R$  there is a *unique*  $b \in R$  such that  $aba = a$ .
  - (a) Show that  $R$  has no [non-zero] zero divisors.
  - (b) Show that for  $a$  and  $b$  as above, we also have  $bab = b$ .
  - (c) Show that  $R$  has 1.
2. Let  $R$  be a commutative ring and  $a \in R$  such that  $a^n \neq 0$  for all positive integers  $n$ . Let  $I$  be an ideal maximal with respect to the property that  $a^n \notin I$  for any positive integer  $n$ . Show that  $I$  is prime.

## Part III

1. Let  $V = \mathbb{R}^2$  and  $\{e_1, e_2\}$  be a basis of  $V$ . Show that  $e_1 \otimes e_2 + e_2 \otimes e_1 \in V \otimes_{\mathbb{R}} V$  cannot be written as a single tensor.
2. Let  $R$  be a PID.
  - (a) Prove that a finitely generated  $R$ -module  $M$  is free if and only if it is torsion free.
  - (b) Prove that if a finitely generated  $R$ -module  $M$  is projective, then it is free.

## Part IV

1. Let  $\mathbb{F}_p$  be the field with  $p$  elements,  $\bar{\mathbb{F}}_p$  be a fixed algebraic closure of  $\mathbb{F}_p$  and let

$$L = \{\alpha \in \bar{\mathbb{F}}_p : p \nmid [\mathbb{F}_p[\alpha] : \mathbb{F}_p]\}.$$

Show that  $L$  is a field.

2. Let  $p$  be a prime,  $F$  be a field of characteristic different from  $p$  and  $f = x^p - a \in F[x]$  [not necessarily irreducible]. Let  $K$  be the splitting field of  $x^p - 1$  over  $F$  and assume that all roots of  $f$  lie in  $K$ .
  - (a) Show that if  $f(\alpha) = 0$  with  $\alpha \notin F$ , then  $F[\alpha] = K$ .
  - (b) Prove that  $f$  has a root in  $F$ .

# ALGEBRA PRELIMINARY EXAM

AUGUST 2012

**Instructions:** Attempt *all* problems in all four parts. Justify each answer.

**General Assumptions:** All rings have  $1 \neq 0$  [and their subrings contain 1] and all modules are unitary.

## Part I

1. Let  $G$  and  $H$  be finite Abelian groups. Prove that if  $G \times H \times H \cong G \times G \times H$ , then  $G \cong H$ .
2. Let  $p$  be a prime and  $G$  be a group of order  $p^n$ . For  $k \in \{1, 2, 3, \dots, (n-1)\}$ , let  $s_k$  and  $n_k$  denote the number of subgroups and normal subgroups of  $G$  of order  $p^k$  respectively. Show that  $s_k - n_k$  is divisible by  $p$ .

## Part II

1. Let  $R$  be a commutative ring for which every proper ideal is prime. Show that  $R$  is a field.
2. Let  $F$  be a field and consider the subring  $R$  of  $F[t]$  given by polynomials with the coefficient of  $t$  equal to zero, i.e.,  $R = F + t^2F[t]$ .
  - (a) Show that  $R$  has an irreducible element which is not prime. [Hence,  $R$  is not PID.]
  - (b) Show that  $R$  is Noetherian. [Hint: Consider a connection between  $R$  and  $F[x, y]$ .]

## Part III

1. Let  $R$  be a commutative ring,  $S$  be a subring of  $R$ ,  $A$  be an  $R$ -module and

$$\mathcal{H} \stackrel{\text{def}}{=} \text{Hom}_R(R \otimes_S (S \oplus S), A).$$

Show that for every *surjective* homomorphism of  $R$ -modules  $\phi : M \rightarrow N$  and  $R$ -module homomorphism  $f : \mathcal{H} \rightarrow N$  there is an  $R$ -module homomorphism  $F : \mathcal{H} \rightarrow M$  such that  $\phi \circ F = f$  if and only if the same is true if we replace  $\mathcal{H}$  by  $A$ .

2. Let  $R$  be a commutative ring,  $D$ ,  $M$  and  $N$  be  $R$ -modules,  $\phi : M \rightarrow N$  be an  $R$ -module homomorphism and  $1 \otimes \phi : D \otimes_R M \rightarrow D \otimes_R N$  be the homomorphism for which

$$(1 \otimes \phi)(d \otimes m) = d \otimes \phi(m).$$

- (a) Assume that  $\phi$  is injective. Show that if  $D$  is free and of finite rank, then  $1 \otimes \phi$  is also injective. [The finite rank is not necessary, but we assume it here for simplicity.]
- (b) Show that the above statement is not true for an arbitrary  $D$ .

## Part IV

1. Let  $F$  be a field and  $K/F$  be an algebraic extension. Show that if  $R$  is a *subring* of  $K$  with  $F \subseteq R \subseteq K$ , then  $R$  is a field.
2. Let  $F$  be a field,  $K/F$  be a Galois extension and  $f \in F[x]$  be monic, separable and irreducible. Show that if  $f = f_1 \cdots f_k$  is the factorization of  $f$  in  $K[x]$ , with  $f_i$  irreducible and monic, then the  $f_i$ 's are distinct, of the same degree and  $G \stackrel{\text{def}}{=} \text{Gal}(K/F)$  acts *transitively* on  $\{f_1, \dots, f_k\}$ . [I.e., given  $\sigma \in G$ , the map  $f_i \mapsto f_i^\sigma$  is a permutation of the  $f_i$ 's and given  $i, j \in \{1, \dots, k\}$ , there is a  $\tau \in G$  such that  $f_i^\tau = f_j$ .]

ALGEBRA PRELIMINARY EXAMINATION  
Spring 2012

Attempt all four parts. Justify your answers.

Part I.

1. Show that a group of order 455 is necessarily cyclic.
2. Let  $G$  be a group of order 56. Show that  $G$  is solvable.

Part II.

1. Let  $f : \mathbb{Q} \rightarrow \mathbb{Z}$  be a function such that  $f(ab) = f(a)f(b)$  for all  $a, b \in \mathbb{Q}$ . Show that the image of  $f$  has at most three elements and there exist an infinite number of such functions whose image has three elements.
2. Let  $R$  be a PID and let  $J$  denote the intersection of all maximal ideals of  $R$ . If  $a^2 - a$  is in  $J$  for all  $a \in R$ , then show that  $R$  has only finitely many maximal ideals.

Part III.

Note: Rings are assumed to be commutative and with  $1 \neq 0$  and modules are assumed to be unitary.

1. Let  $R$  be an integral domain and let  $M, N$  be projective  $R$ -modules. Show that  $M \otimes_R N$  is a projective  $R$ -module.
2. Suppose  $R$  is a principal ideal domain that is not a field. Suppose  $M$  is a finitely generated  $R$ -module such that for every maximal ideal  $P$  of  $R$ ,  $M/PM$  is a cyclic  $R/P$ -module. Show that  $M$  itself is cyclic.

Part IV.

1. Let  $f(X)$  be a monic polynomial of degree 9 having rational coefficients. Assume that  $f(X)$  is irreducible in  $\mathbb{Q}[X]$ . Let  $K$  denote the splitting field of  $f$  over  $\mathbb{Q}$  and let  $u \in K$  be a root of  $f$ . If  $[K : \mathbb{Q}] = 27$ , then show that  $\mathbb{Q}[u]$  has a subfield  $L$  with  $[L : \mathbb{Q}] = 3$ .
2. Let  $F, K$  be fields such that  $K$  is a finite Galois extension of  $F$  with Galois group  $G$ . Suppose  $a \in K$  is such that  $\sigma(a) - a \in F$  for all  $\sigma \in G$ . If the characteristic of  $F$  does not divide the order of  $G$ , then show that  $a \in F$ . Assuming  $F$  to be the field of two elements, construct a quadratic field extension  $K := F[a]$  of  $F$  such that  $\sigma(a) - a \in F$  for all  $\sigma \in G$ .

## Algebra Preliminary Exam

January 2011

Attempt all problems and justify all your answers.

All rings have an identity  $1 \neq 0$ , all ring homomorphisms send 1 to 1, and all R-modules are unitary.

- I. 1. Let  $G$  be a finite simple group. Show that if  $G$  has a subgroup  $H$  with  $[G:H] = n \geq 2$ , then  $|H| \mid (n-1)!$ .
2. List, up to isomorphism, all groups of order 153. Justify your answer.
- II. 1. Let  $R$  be a commutative ring and  $I$  an ideal of  $R$ . Let  $I^* = (I, X)$  be an ideal of the polynomial ring  $R[X]$ . Determine, in terms of  $I$ , when  $I^*$  is a prime ideal of  $R[X]$  and when  $I^*$  is a maximal ideal of  $R[X]$ . Justify your answers.
2. (a) Show that if a commutative ring  $R$  satisfies DCC on ideals (i.e.,  $R$  is Artinian), then  $R$  has only a finite number of maximal ideals.
- (b) Give an example to show that (a) may be false if DCC is replaced by ACC (i.e., if  $R$  is Noetherian).

**III.** 1. Let  $f: M \rightarrow M$  be an  $R$ -module homomorphism with  $f \circ f = f$ .

Show that the following statements are equivalent.

(a)  $f$  is injective.

(b)  $f$  is surjective.

(c)  $f = 1_M$ .

2. (a) Let  $G$  and  $H$  be finitely generated abelian groups such that  $\mathbb{Z}_n \otimes G \cong \mathbb{Z}_n \otimes H$  for every integer  $n \geq 2$ . Show that  $G \cong H$ .

(b) Give an example to show that (a) may be false if  $G$  and  $H$  are not both finitely generated.

**IV.** 1. Let  $F$  be a subfield of a field  $L$ . Show that  $L/F$  is an algebraic extension if and only if every subring  $R$  of  $L$  containing  $F$  is a field.

2. Compute the Galois group of  $f(X) = X^4 + X + 1 \in \mathbb{Z}_2[X]$ .

ALGEBRA PRELIMINARY EXAMINATION  
Fall 2011

Attempt all four parts. Justify your answers.

Part I.

1. How many Sylow 2-subgroups does  $S_5$  (the group of permutations of  $\{1, 2, 3, 4, 5\}$ ) have ?
2. Let  $G$  be a group of order 231. Show that  $G$  is Abelian if and only if  $G$  has an Abelian subgroup of order 21.

Part II.

Note: Rings are assumed to be commutative and with  $1 \neq 0$ , subrings are assumed to contain 1 and ring-homomorphisms are assumed to map 1 to 1.

1. Let  $R$  be a UFD such that each maximal ideal of  $R$  is a principal ideal. Prove that  $R$  is a PID.
2. Let  $\mathbb{R}[[X]]$  denote the power-series ring in a single indeterminate  $X$  over the field of real numbers  $\mathbb{R}$ . If  $T$  is a multiplicative subset of  $\mathbb{R}[[X]]$  containing 1 but not containing 0, then show that either  $T^{-1}\mathbb{R}[[X]] = \mathbb{R}[[X]]$  or  $T^{-1}\mathbb{R}[[X]]$  is a field.

Part III.

Note: Rings are assumed to be commutative and with  $1 \neq 0$  and modules are assumed to be unitary.

1. Let  $R$  be an integral domain and  $I$  an ideal of  $R$ . Show that there exists a surjective  $R$ -module homomorphism  $f : I \rightarrow R$  if and only if  $I$  is a nonzero principal ideal.
2. Let  $K$  be a field,  $X$  an indeterminate over  $K$  and  $M$  a finitely generated  $K[X]$ -module. Show that  $M$  is a projective  $K[X]$ -module if and only if  $M$  is  $K[X]$ -module isomorphic to  $K[X] \otimes_K V$  for some finite dimensional  $K$ -vector space  $V$ .

Part IV.

1. Let  $K$  be a field and  $F$  a subfield of  $K$ . The group of units of  $K$  is denoted by  $K^\times$ . Suppose  $f \in F[X]$  is a monic irreducible polynomial and  $a, b \in K^\times$  are such that  $f(a) = 0 = f(b)$ . Show that the subgroup of  $K^\times$  generated by  $a$ , is isomorphic to the subgroup of  $K^\times$  generated by  $b$ .
2. Let  $f \in \mathbb{Q}[X]$  be a polynomial of degree 4 such that the Galois group of  $f$  (over  $\mathbb{Q}$ ) is a group of order 6. Show that  $f$  has a root in  $\mathbb{Q}$ .



Attempt all problems and justify all answers. All rings have an identity  $1 \neq 0$ , ring homomorphisms send 1 to 1, and all  $R$ -modules are unitary.

- I.** 1. Let  $f : G \rightarrow H$  be a surjective homomorphism of finite groups and  $y \in H$  with  $|y| = n$ . Show that there is an  $x \in G$  with  $|x| = n$ .
2. Let  $p$  and  $q$  be primes,  $p \geq q$ ,  $n \geq 1$ , and  $G$  a group with  $|G| = p^n q$ . Show that  $G$  has a normal subgroup  $H$  of order  $p^n$ . (Hint: do the  $p > q$  and  $p = q$  cases separately.)
- II.** 1. Let  $R$  be a commutative ring with distinct prime ideals  $P$  and  $Q$  with  $P \cap Q = \{0\}$ . Show that  $R$  is isomorphic to a subring of the direct product of two fields.
2. Let  $p$  and  $q$  be positive primes. Show that the polynomial  $f(X) = X^3 + pX^2 + q \in \mathbb{Z}[X]$  is irreducible in  $\mathbb{Q}[X]$ .
- III.1.** Let  $A$  and  $B$  be finite abelian groups with  $|A| = m$  and  $|B| = n$ . Show that  $\text{Hom}_{\mathbb{Z}}(A, B) = 0$  if and only if  $\text{gcd}(m, n) = 1$ .
2. Let  $A$  be a submodule of a projective  $R$ -module  $B$ . Show that  $A$  is projective if  $B/A$  is projective.

- IV.** 1. Let  $K \subseteq F$  and  $K \subseteq L$  be subfields of a field  $M$  with  $[F:K] = p$  and  $[L:K] = q$  for distinct primes  $p$  and  $q$ . Show that  $F \cap L = K$ , and that  $F = K(\alpha)$  and  $L = K(\beta)$  for any  $\alpha \in F - K$  and  $\beta \in L - K$ .
2. Let  $K$  be a field and  $f(X) \in K[X]$  be irreducible and separable with  $\deg(f(X)) = n$ . Show that if the Galois group  $G$  of  $f(X)$  over  $K$  is abelian, then  $|G| = n$ .